

## Angiotensin II receptor antagonists against migraine in pregnancy: fatal outcome

Kirsti Haaland

Received: 26 September 2009 / Accepted: 9 December 2009 / Published online: 9 January 2010  
© Springer-Verlag 2010

**Abstract** A pregnant young woman with a severe migraine is prescribed candesartan, an angiotensin II type 1 receptor antagonist (AT II antagonists). This has a positive effect—except for severe maldevelopment of her fetus. There is an increase in the use of the fetotoxic drugs, AT II antagonists and angiotensin-converting enzyme inhibitors, as prophylactic treatment of migraines, in addition to their use as hypertensives.

**Keywords** Migraine · Pregnancy · Fetotoxic medication · Angiotensin II receptor antagonist · Angiotensin-converting enzyme inhibitor · Hypertension

### Abbreviations

AT II antagonist    Angiotensin II receptor antagonists  
ACE-inhibitors    Angiotensin-converting enzyme inhibitors

### Introduction

There is an increase in the use of angiotensin II type 1 receptor antagonists (AT II antagonists) and angiotensin-converting enzyme inhibitors (ACE-inhibitors) as prophylactic treatment of migraines, in addition to their use as antihypertensives. These drugs pass the placenta and have highly adverse effects on the renal organogenesis, leading

to maldevelopment firstly of the renal system and secondarily of the lungs [1, 2].

### Case story

A 35-year-old woman has a missed abortion. She is given hospital treatment. The fetus is autopsied without conclusive diagnosis. In the woman's next pregnancy, only 6 weeks later, an ultrasound examination is conducted in week 12 and again in week 30. One week later, the midwife finds a declining symphysis to fundus increment, and the woman is therefore immediately referred to the hospital. Anhydramnion is diagnosed. There is severe pathology of the kidneys and bladder. The development of the skull, thorax and possibly the intestines are also pathological. The reasons for the maldevelopment are unclear.

The woman is healthy, apart from suffering a severe common migraine. Before pregnancy, she had three to four attacks every week of unilateral, throbbing pain associated with nausea, vomiting and photophobia, lasting 6–48 h, preceded by transient scintillating scotomas. During pregnancy, the attacks are milder, shorter and less frequent. Throughout both pregnancies, she has been taking candesartan (16 mg/day), pramipexole (0.18 mg 3×) and amitriptyline (25 mg/day) as prophylaxis against migraines, in addition to zolmitriptan and metoclopramide during attacks. Candesartan treatment was initiated more than a year before her first pregnancy by an experienced neurologist who was also consulted during pregnancy. The medication was known to her general practitioner and the doctors who conducted the ultrasound examinations. The hospital obstetricians documented the medication in the admission notes in both pregnancies.

---

K. Haaland (✉)  
Department of Pediatrics, Oslo University Hospital,  
Ullevål, Kirkev 166, 0407 Oslo, Norway  
e-mail: kirsti66@online.no; kirsti.haaland@ulleva.no

Nonetheless, the fetus is 33 weeks before doctors become aware that the medication is fetotoxic. Candesartan, an AT II antagonist, is seponated. When the baby is born, he has renal tubular dysgenesis, hypoplasia of the skull and the lungs, and hyaline membranes of the lungs. This is not compatible with life.

## Materials and methods

MEDLINE and the Norwegian Database of Adverse Effects [3] were searched for descriptions of fetal injuries related to AT II antagonists ACE-inhibitors. Information about the extent of use of these drugs in Norway was obtained from the National Prescription Database.

## Results

Angiotensin II receptor antagonists and ACE-inhibitors reduce blood pressure by blocking the renin-angiotensin system. They also have a positive, prophylactic effect against migraines, although the mechanisms are poorly understood. The colocalization of AT1, glutamate and GABA receptors on medullary rostral ventromedial neurons suggests a nociceptive modulatory [4]. Unfortunately, these drugs cross the placenta and affect the circulation of the fetal kidneys, and, more importantly, reduce stimulation of AT II receptors. This has a highly adverse effect on the renal organogenesis in the second and third trimesters [5]. The kidneys develop abnormally and are unable to produce urine; there is oligohydramnion and thereby, inter alia, maldevelopment of the lungs [1, 2].

The summary of product characteristics in the technical brochures clearly states that these drugs are fetotoxic and should not be used during pregnancy [6]. More than 20 cases of fetal injury/maldevelopment after exposure to AT II antagonists and ACE-inhibitors have been reported in the literature [1, 2]. Two cases have been reported to the Norwegian Database of Adverse Effects. Norway has 4.7 million inhabitants.

In Norway, 1.2% of women aged 30–39 years were dispensed drugs in 2007 that affect the renin-angiotensin system, an increase compared to 2004. More than 50% of childbearing women in Norway are more than 30 years old. The mean age at delivery is increasing and will thus cause the number of pregnant women with hypertension requiring medical treatment to increase. In addition, there is reason to believe that there is an increase in the use of these drugs against migraines. They are well tolerated, and several studies have demonstrated their positive prophylactic effects [7–11]. In American and European guidelines, candesartan is listed among second- and third-line agents,

respectively, for migraine prophylaxis [12–14]. In Australia it is not yet listed as an appropriate agent, but is widely used by the neurologists [15]. This also appears to be the case in Norway, although it is difficult to document, as these are not approved drugs against migraines in this country.

## Discussion

Migraine is a most disabling disorder, which can be extremely difficult to treat. A variety of drugs from diverse pharmacological classes are in use for migraine prevention. AT II antagonists and ACE-inhibitors are traditional anti-hypertensives that have proved to be effective also in migraine prophylaxis. Their fetotoxic effects have been demonstrated in humans [1, 2] and well documented in animal research. When administered to rats, mice or piglets during renal development these drugs induce severe renal histological abnormalities, including papillary atrophy, tubulointerstitial fibrosis and tubular atrophy and dilatation [16, 17]. Nonetheless, there is reason to believe that an increasing number of women of reproductive age will use these drugs. They should be advised of the possible hazards, and treatment should be stopped as soon as pregnancy is planned or detected.

**Conflict of interest** None.

## References

1. Kato K, Okuda M, Ishikawa H, Takahashi T, Hirahara F (2008) Oligohydramnios and pulmonary hypoplasia: a case in which involvement of an angiotensin II receptor antagonist was suspected. *J Obstet Gynaecol Res* 34(2):242–246
2. Serreau R, Luton D, Macher MA, Delezoide AL, Garel C, Jacqz-Aigrain E (2005) Developmental toxicity of the angiotensin II type 1 receptor antagonists during human pregnancy: a report of ten cases. *BJOG* 112(6):710–712
3. Regional medicines information centres. Norwegian database of adverse effects
4. Ramadan NM (2007) Current trends in migraine prophylaxis. *Headache* 47(Suppl 1):S52–S57
5. Lacoste M, Cai Y, Guicharnaud L et al (2006) Renal tubular dysgenesis, a not uncommon autosomal recessive disorder leading to oligohydramnios: role of the renin–angiotensin system. *J Am Soc Nephrol* 17(8):2253–2263
6. Norwegian Medicines Agency. Summary of product characteristics (SPC), Atacand
7. Tronvik E, Stovner LJ, Helde G, Sand T, Bovim G (2003) Prophylactic treatment of migraine with an angiotensin II receptor blocker: a randomized controlled trial. *JAMA* 289(1):65–69
8. Schrader H, Stovner LJ, Helde G, Sand T, Bovim G (2001) Prophylactic treatment of migraine with angiotensin converting enzyme inhibitor (lisinopril): randomised, placebo controlled, crossover study. *BMJ* 322(7277):19–22
9. Etminan M, Levine MA, Tomlinson G, Rochon PA (2002) Efficacy of angiotensin II receptor antagonists in preventing

- headache: a systematic overview and meta-analysis. *Am J Med* 112(8):642–646
10. Bender WI (1995) ACE inhibitors for prophylaxis of migraine headaches. *Headache* 35(8):470–471
  11. Owada K (2004) Efficacy of candesartan in the treatment of migraine in hypertensive patients. *Hypertens Res* 27(6):441–446
  12. Silberstein SD (2000) Practice parameter: evidence-based guidelines for migraine headache (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 55(6):754–762
  13. Evers S, Afra J, Frese A et al (2006) EFNS guideline on the drug treatment of migraine—report of an EFNS task force. *Eur J Neurol* 13(6):560–572
  14. Modi S, Lowder DM (2006) Medications for migraine prophylaxis. *Am Fam Phys* 73(1):72–78
  15. Stark RJ, Stark CD (2008) Migraine prophylaxis. *Med J Aust* 189(5):283–288
  16. Sanchez SI, Seltzer AM, Fuentes LB, Forneris ML, Ciuffo GM (2008) Inhibition of Angiotensin II receptors during pregnancy induces malformations in developing rat kidney. *Eur J Pharmacol* 588(1):114–123
  17. Guron G, Sundelien B, Wickman A, Friberg P (1988) Angiotensin-converting enzyme inhibition in piglets induces persistent renal abnormalities. *Clin Exp Pharmacol Physiol* 25(2):88–91