

POSTER PRESENTATION

Open Access

# The differences in trigeminal system functional state in chronic and frequent migraine patients

G Pavlic<sup>1\*</sup>, S Odobescu<sup>1</sup>, I Moldovanu<sup>1</sup>, V Lacusta<sup>2</sup>, L Rotaru<sup>1</sup>, G Corcea<sup>1</sup>

From The European Headache and Migraine Trust International Congress  
London, UK. 20-23 September 2012

## Introduction

Electrophysiological tools like trigeminal somato-sensory evoked potentials (TSSEP) are used for evaluation of trigeminal system (TS) functional state. The activity of the later is an important factor in migraine pathogenesis. The study aim was to analyze the TS functional state in chronic and frequent migraine patients in relation with age during the migraine attack (critic period) and in the interictal period on lateralized stimulation (left/right).

## Results

We have included 170 women with migraine (according to ICHD-II, 2006 criteria), between 15-55 years, 109-chronic migraine (CM) and 61-frequent migraine (FM), mean age  $38.3 \pm 10.7$  and  $36.5 \pm 9.1$  years, respectively. In CM vs. FM patients aged 15-25 years a statistic significant reduction in wave latencies N15 ( $15.4 \pm 0.47$  vs.  $17.67 \pm 0.42$ ,  $p < 0.01$ ), P22 ( $22.24 \pm 0.69$  vs.  $25.11 \pm 0.73$ ,  $p < 0.05$ ), interval N6-P9 ( $3.56 \pm 0.29$  vs.  $5.11 \pm 0.56$ ,  $p < 0.05$ ) on the left side and of intervals N6-P9 ( $3.28 \pm 0.40$  vs.  $5.36 \pm 0.54$ ,  $p < 0.05$ ) and N6-P22 ( $16.76 \pm 0.52$  vs.  $19.18 \pm 0.80$ ,  $p < 0.05$ ) on the right was observed, but not in other age groups. Reduced latency and shorter conduction time at nuclear-thalamic level (interval N6-P9) might reflect an increased excitability of thalamic and cortical structures (P22) of trigeminal sensory complex in young patients. There were no differences in TSSEP in CM vs. FM during a migraine attack.

## Conclusions

Functional disturbances in trigeminal system in chronic migraine vs. frequent migraine might be seen in young patients based on latencies of TSSEP and on central time of the somatosensory trigeminal transmission. These changes

might reflect a different activity of brainstem-diencephalic (thalamic) structures.

## Author details

<sup>1</sup>Institute of Neurology and Neurosurgery, Moldova, Republic of. <sup>2</sup>State Medical and Pharmaceutical University, Moldova, Republic of.

Published: 21 February 2013

## Reference

1. Ambrosini A, de Noordhout AM, Sándor PS, Schoenen J: Electrophysiological studies in migraine: a comprehensive review of their interest and limitations. *Cephalalgia* 2003, **23**(Suppl. 1):13-31.

doi:10.1186/1129-2377-14-S1-P125

Cite this article as: Pavlic et al.: The differences in trigeminal system functional state in chronic and frequent migraine patients. *The Journal of Headache and Pain* 2013 **14**(Suppl 1):P125.

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)

<sup>1</sup>Institute of Neurology and Neurosurgery, Moldova, Republic of  
Full list of author information is available at the end of the article