

POSTER PRESENTATION

Open Access

# Monitoring cortical neuronal activity and spreading depression in freely behaving familial hemiplegic migraine *Cacna1a* R192Q knockin mice

T Houben<sup>1</sup>, R Shyti<sup>2</sup>, Q Dees<sup>2</sup>, S van Berloo<sup>3</sup>, L de Groot<sup>1</sup>, GM Terwindt<sup>1</sup>, MD Ferrari<sup>1</sup>, EA Tolner<sup>1\*</sup>, AM van den Maagdenberg<sup>2</sup>

From The European Headache and Migraine Trust International Congress London, UK. 20-23 September 2012

## Introduction

Experimental findings from transgenic migraine mouse models that carry a human FHM1 gain-of-function mutation in CaV2.1 (P/Q-type) calcium channels underscore the role of neuronal hyperexcitability in migraine [1,2]. However, functional data that link the excitability changes to neuronal network activity and the enhanced propensity to cortical spreading depression (CSD), the likely mechanism underlying migraine aura, are largely lacking.

## Purpose/background/objectives

Here, we aimed to set up an electrophysiology platform to study changes in cortical neuronal network activity in relation to CSD in freely behaving transgenic migraine mice.

## Methods

We developed an electrophysiology system for long-term recordings of DC-EEG and multi-unit-activity from the cortex of freely behaving mice. The system combines a counterbalanced 7 channel swivel with custom-built differential DC-EEG, AC-EEG and unit activity amplifiers. Stable DC-EEG recordings are obtained using AgAgCl epidural electrodes, while intracortical platinum electrodes are used for simultaneous recording of multi-unit-activity and AC-EEG. For CSD induction intracortical microdialysis was used for infusion of high KCl solution.

## Results

Simultaneous recordings of multi-unit activity, DC- and AC-EEG were made from the sensorimotor and occipital

cortex of wild-type and FHM1 migraine mice for up to 3 weeks. Apart from spontaneous cortical activity, visual evoked cortical responses were induced using 1 ms blue light pulses. Microdialysis with KCl solution resulted in successful induction of CSD events in the awake mice.

## Conclusion

We established a novel platform for performing longitudinal recordings of cortical neuronal activity and spreading depression in freely behaving mice carrying migraine-specific mutations. Using this platform, we aim to characterize how cortical activity is altered by modulatory factors that predispose for migraine attacks.

## Author details

<sup>1</sup>Department of Neurology, Leiden University Medical Center, Leiden, Netherlands. <sup>2</sup>Department of Human Genetics, Leiden University Medical Center, Leiden, Netherlands. <sup>3</sup>Instrument Development Department, Leiden University Medical Center, Leiden, Netherlands.

Published: 21 February 2013

## References

1. van den Maagdenberg AMJM, Pietrobon D, Pizzorusso T, Kaja S, Broos LAM, Cesetti T, et al: A *Cacna1a* knockin migraine mouse model with increased susceptibility to cortical spreading depression. *Neuron* 2004, **41**(5):701-710.
2. Tottene A, Conti R, Fabbro A, Vecchia D, Shapovalova M, Santello M, et al: Enhanced excitatory transmission at cortical synapses as the basis for facilitated spreading depression in Ca(v)2.1 knockin migraine mice. *Neuron* 2009, **61**(5):762-773.

doi:10.1186/1129-2377-14-S1-P79

**Cite this article as:** Houben et al.: Monitoring cortical neuronal activity and spreading depression in freely behaving familial hemiplegic migraine *Cacna1a* R192Q knockin mice. *The Journal of Headache and Pain* 2013 **14**(Suppl 1):P79.

<sup>1</sup>Department of Neurology, Leiden University Medical Center, Leiden, Netherlands

Full list of author information is available at the end of the article