

MEETING ABSTRACT

Open Access

EHMTI-0230. Characteristics of menstrual and nonmenstrual migraine attacks in women with menstrual migraine

KG Vetvik^{1*}, EA MacGregor², J Šaltyté Benth³, AC Lundqvist⁴, MB Russell⁵

From 4th European Headache and Migraine Trust International Congress: EHMTIC 2014 Copenhagen, Denmark. 18-21 September 2014

Introduction

The International Classification of Headache Disorders 3 β defines menstrual migraine (MM) as attacks of migraine without aura occurring on day 1 \pm 2 of the menstrual cycle in \geq 2/3 menstruations. Previous studies show conflicting results regarding possible differences between menstrual and nonmenstrual migraine attacks.

Aim

To compare characteristics of menstrual and nonmenstrual migraine attacks in women prospectively diagnosed with MM.

Method

237 women from the general population with self-reported migraine in \geq 50% of their menstruations were interviewed and diagnosed by a neurologist. Subsequently they were asked to complete a three month prospective headache- and menstruation diary. When a headache occurred, the women recorded information about pain intensity, quality and location, time of onset and end of headache, associated symptoms, aggravation by routine physical activity, medical treatment and sick leave. In addition, they were asked to record each day of uterine bleeding.

Results

Of the 123 (52%) returned diaries, 42 were excluded due to oligoamenorrhoea (n = 36), no migraine (n = 4) or incompleteness (n = 2). Among the remaining 81 women, 56 fulfilled the diagnostic criteria for MM. In these

56 women, menstrual migraine attacks were more painful on a 0-10 scale (coefficient 0.34, 95% CI 0.02-0.65), longer lasting (coefficient 10.91 hours, 95% CI 5.68-16.14), and required more doses of symptomatic treatment (coefficient 1.09 doses, 95% CI 0.50-1.68) compared to nonmenstrual attacks. Additionally, they were more often associated with nausea (OR 2.22, 95% CI 1.35-3.65).

Conclusion

In women with MM, menstrual migraine attacks are more severe than nonmenstrual attacks.

No conflict of interest.

Authors' details

¹Head and Neck Research Group Research Centre and Institute of Clinical Medicine, University of Oslo and Akershus University Hospital, Lørenskog, Norway. ²Centre for Neuroscience and Trauma Blizard Institute, Barts and the London School of Medicine and Dentistry, London, UK. ³Institute of Clinical Medicine and HØKH Research Centre, University of Oslo and Akershus University Hospital, Lørenskog, Norway. ⁴Institute of Clinical Medicine and Head and Neck Research Group and HØKH Research Centre and Department of Neurology, University of Oslo and Akershus University Hospital, Lørenskog, Norway. ⁵Institute of Clinical Medicine and Head and Neck Research Group Research Centre, University of Oslo and Akershus University Hospital, Lørenskog, Norway.

Published: 18 September 2014

doi:10.1186/1129-2377-15-S1-D67

Cite this article as: Vetvik et al.: EHMTI-0230. Characteristics of menstrual and nonmenstrual migraine attacks in women with menstrual migraine. *The Journal of Headache and Pain* 2014 **15** (Suppl 1):D67.

¹Head and Neck Research Group Research Centre and Institute of Clinical Medicine, University of Oslo and Akershus University Hospital, Lørenskog, Norway

Full list of author information is available at the end of the article