Open Access

EHMTI-0228. A clinical interview versus prospective headache diaries in the diagnosis of menstrual migraine without aura

KG Vetvik^{1*}, EA MacGregor², AC Lundqvist³, MB Russell⁴

From 4th European Headache and Migraine Trust International Congress: EHMTIC 2014 Copenhagen, Denmark. 18-21 September 2014

Introduction

The International Classification of Headache Disorders (ICHD) II and III beta defines menstrual migraine as attacks of migraine without aura occurring on day 1 ± 2 of the menstrual cycle in $\geq 2/3$ menstruations. According to the ICHD III beta, a three month prospective headache diary is required in order to establish the diagnosis of MM.

Aims

To compare the diagnosis of MM from a clinical interview based on the ICHD II-criteria for MM, with prospective headache diaries in a population-based study.

Methods

237 women with self-reported migraine in at least half of menstruations were interviewed by a neurologist about headache and diagnosed according to ICHD II. Additionally, the MM-criteria were expanded to include other types of migraine related to menstruation. Subsequently, all women were asked to complete three month prospective headache diaries.

Results

A total of 123 (52%) women completed both clinical interview and diaries. Thirty-eight women were excluded from the analyses; two had incomplete diaries and 36 women recorded £1 menstruation, leaving 85 diaries eligible for analysis. Sensitivity, specificity, positive and negative predictive value and Kappa for the diagnosis of MM in clinical interview vs. headache diary were 82%, 83%, 90%, 71% and 0.62 (95% CI 0.46-0.80). Using

¹Institute of Clinical Medicine and Head and Neck Research group Research Centre, University of Oslo and Akershus University Hospital, Lorenskog, Norway

Full list of author information is available at the end of the article



a broader definition of MM, Kappa was 0.64 (95% CI 0.47-0.83).

Conclusion

A thorough clinical interview is valid for the diagnosis of MM. When this is undertaken, prospective headache diaries should not be mandatory to diagnose MM but may be necessary to exclude a chance association.

No conflict of interest.

Authors' details

¹Institute of Clinical Medicine and Head and Neck Research group Research Centre, University of Oslo and Akershus University Hospital, Lorenskog, Norway. ²Barts and the London School of Medicine and Dentistry, Centre for Neuroscience and Trauma Blizard Institute, London, UK. ³Head and Neck Research Group and HØKH Research Centre and Institute of Clinical Medicine and Department of Neurology, Akershus University Hospital and University of Oslo, Lørenskog, Norway. ⁴Head and Neck Research Group Research Centre and Institute of Clinical Medicine, Akershus University Hospital and University of Oslo, Lørenskog, Norway.

Published: 18 September 2014

doi:10.1186/1129-2377-15-S1-D68

Cite this article as: Vetvik *et al.*: EHMTI-0228. A clinical interview versus prospective headache diaries in the diagnosis of menstrual migraine without aura. *The Journal of Headache and Pain* 2014 **15**(Suppl 1):D68.

