

Psychiatric comorbidity in patients with chronic daily headache and migraine: a selective overview including personality traits and suicide risk

Maurizio Pompili · Daniela Di Cosimo ·
Marco Innamorati · David Lester ·
Roberto Tatarelli · Paolo Martelletti

Received: 17 April 2009 / Accepted: 19 May 2009 / Published online: 25 June 2009
© Springer-Verlag 2009

Abstract Studies on the prevalence and impact of psychiatric disorders among headache patients have yielded findings that have clarified the relationship between migraine and major affective disorders, anxiety, illicit drug abuse, nicotine dependence, and suicide attempts. Studies in both clinical and community-based settings have demonstrated an association between migraine and a number of specific psychiatric disorders. In large-scale population-based studies, persons with migraine are from 2.2 to 4.0 times more likely to have depression. In longitudinal studies, the evidence supports a bidirectional relationship between migraine and depression, with each disorder increasing the risk of the other disorder. Although a strong association has been demonstrated consistently for migraine and major depression, especially for migraine with aura,

there has been less systematic research on the links between migraine and bipolar disorder. This review will focus on the way in which psychiatric disorders decrease the quality of life and result in a worse prognosis, chronicity of the disease, and a worse response to treatment. Short-term pharmaceutical care intervention improves the patients' mental health, but it does not significantly change the number and severity of headaches. The increase in self-efficacy and mental health associated with pharmaceutical care may be instrumental in improving the long-term pharmacotherapy of patients with migraine and headache.

Keywords Psychiatric disorders · Headache · Suicide risk · Treatment

M. Pompili (✉) · D. Di Cosimo · R. Tatarelli
Department of Psychiatry, Sant'Andrea Hospital,
"Sapienza" University of Rome,
1035 Via di Grottarossa, 00189 Rome, Italy
e-mail: maurizio.pompili@uniroma1.it;
mpompili@mclean.harvard.edu

M. Pompili
McLean Hospital Harvard Medical School,
Belmont, MA, USA

M. Innamorati
Università Europea di Roma, Rome, Italy

D. Lester
The Richard Stockton College of New Jersey,
Pomona, NJ, USA

P. Martelletti
Department of Medical and Molecular Sciences,
2nd School of Medicine, Sant'Andrea Hospital,
"Sapienza" University of Rome, Rome, Italy

Introduction

Chronic daily headache (CDH) refers to a group of disorders in which headache occurs 15 or more days per month for at least 3 months [1–5] and includes: chronic migraine (CM), migraine with or without aura and migraine aura without headache, chronic tension-type headache (CTTH), hemicrania continua (HC), and new daily persistent headache [6–8].

Chronic migraine (CM) affects 2.4% of the general population [5] and it is often comorbid with other conditions, such as depression (80%) [9], anxiety (70%) [9, 10], insomnia (71%) [11], chronic fatigue (66%) [12], and fibromyalgia (35%) [13]. Recently, studies have viewed the relationship between migraine and psychiatric disorder as (1) bi-directional [14] since migraine patients have a more than threefold risk of developing depression and (2) specific [15] since the presence of migraine increases a patient's risk of developing depression or panic attack

disorder and the presence of depression or panic attack disorder is associated with a greater risk of developing migraine.

Sheftel and Atlas [16], looking at clinical manifestations associated with daily chronic headache, found a high incidence of anhedonia, decreased concentration, decreased libido, decreased energy, and sleep disorders, all characteristic elements of depression. They proposed that headache is not simply a symptom of depression, but that headache and several mood and anxiety disorders are clinical entities that share some pathophysiological bases [15, 17–24], most likely disturbances in the serotonergic systems [20, 25–27].

Epidemiology

Epidemiological research has indicated an association between migraine and mood disorder with a lifetime prevalence of major depression three times higher among patients with migraine when compared with patients without migraine [28]. This comorbidity between CDH and psychiatric disorders seems to be more frequent in women, in chronic forms of migraine [29] and in first-degree relatives of patients with mood disorder and migraine [21]. Migraine with aura is strongly associated with an early age of onset of the mood disorder (OR 11.4, age < 15 years; OR 5.6, age 15–24 years) [20]. Migraine, weekly or daily, occurs in 3–15% of children [30, 31] while 9–33% of individuals suffer from non-migraine headache at least monthly [31]. The prevalence of migraine is highest among patients with bipolar II disorder (77%), and this is significantly different from the prevalence among bipolar I patients (14%) [32]. The unipolar group has a prevalence of 46%, in between the two bipolar groups [32]. This is in broad agreement with the study by Endicott [33], who found a 51% prevalence of migraine in patients with characteristics similar to bipolar II

patients. The results of some epidemiological studies are shown in Table 1.

Psychiatric disorders and headache

The presence of migraine or severe non-migraine headache increases a patient's risk of developing depression or panic attack disorder, whereas the presence of depression or panic attack disorder is associated with a greater risk of developing migraine, but not severe non-migraine headache [15]. Merikangas and colleagues [17], in a prospective study, supported the idea that anxiety induces or facilitates the onset of a primary headache, acting as trigger for the subsequent development of a mood disorder such as depression. People suffering from migraine were 12 times more likely to suffer from panic attacks [21] and 2.5 times more likely to suffer from depression.

Psychiatric comorbidity is more frequent in chronic pain syndromes, such as the chronic form of migraine [34]. It is important to distinguish between migraine with or without aura and aura without headache. In one population-based epidemiological study [17, 35], migraine with aura was associated with anxiety disorders, recurrent brief depression and hypomania, whereas only the phobic and panic disorders were more frequent among patients suffering from migraine without aura. Patients with tension-type headache did not differ from controls with respect to any of the affective or anxiety disorders. Oedegaard and colleagues [20] showed that a large number of patients presenting with major affective disorders have migraine aura without headache. These patients had a lower frequency of an affective temperament and a lower probability of having made a suicide attempt, as well as a higher age of onset of migraine auras relative to patients suffering from migraine with aura.

The more common psychiatric conditions related to migraine are depression, bipolar disorders, anxiety [(especially,

Table 1 Prevalence of migraine in bipolar disorder from previous studies

Source	Method	Sample size	Migraine prevalence (%)
Blehar et al. (1998)	Medical section of DIGS	<i>N</i> = 327	Total 21.1
		<i>F</i> = 186	Females 26.5
Cassidy and Flanagan (1957)	Self-report of headache	<i>N</i> = 100	Total 49.0
Mahmood et al. (1999)	Self-report of IHS migraine criteria	<i>N</i> = 81	Total 25.9
		<i>F</i> = 37	Females 27.0
Marchesi et al. (1989)	Diagnosis by neurologist	<i>N</i> = 30	Total 20.0
Fasmer (2001)	Administered IHS migraine criteria	<i>N</i> = 27	Bipolar I 13.0 Bipolar II 77.0
Younes et al. (1986)	Mother's report of past diagnosis of migraine	<i>N</i> = 21	Total 28.6
			Females = 0
			Male = 21

DIGS Diagnostic Interview for Genetic Studies, IHS International Headache Society

generalized anxiety disorders (GAD)] [17, 36] and somatoform disorders. These comorbid disorders have been identified in several epidemiological studies [22, 37–40] as well as in clinical studies of treatment-seeking patients [34, 40, 41]. The association between migraine and depression has also been studied in a number of large-scale community studies (Table 2).

The association of CDH with psychiatric disorders is controversial, especially with regard to depression (Table 3). Verri and colleagues [10] found an association between CDH and at least one psychiatric disorder in 90% of their patients, mainly GAD, followed by major depression (25%) and dysthymia (17%). Verri [10] and later Juang [48] confirmed that a longer major depressive disorder (or chronic depression) was the most frequent comorbidity among these patients, especially when the chronic headache had lasted for more than 5 years. However, Juang and colleagues [48] found that the frequency of any type of anxiety disorder was significantly higher in patients with chronic migraine than in those with CTTH. In prior research, the comorbidity between tension-

type headache and psychiatric disorders has been investigated only in clinical populations, in which it has been showed that this association is more common in the chronic form of tension-type headache [14]. In fact, the frequency of anxiety and mood disorders were 3–15 times higher in CTTH patients than in controls [14], and the frequencies reported were 21% for depression and 17% for anxiety disorders in these patients [41], with significantly higher anxiety and depression scores [41, 49]. Moreover, in previous studies, especially in the chronic form of tension-type headache, affective disorders were found to be frequent [50, 51]. Of the three studies conducted in clinical samples, none reported any significant differences in psychiatric comorbidity between migraine and tension-type headache [49, 52, 53].

There has been less research on the role of personality disorders in headache. Mathew [54] found significantly higher abnormal personality profiles in CDH patients than in controls using the Minnesota Multiphasic Personality Inventory (MMPI) but found no difference in the profiles of patients with different subtypes of CDH. In a small sample

Table 2 Select studies of the association between migraine and depression

Study	Method	Migraine and depression OR (95% CI)	Bi-directional relationship OR (95% CI)
Longitudinal studies			
Breslau et al. [37]	IHS migraine criteria	Not assessed	New-onset migraine 3.5 (2.2–5.6) New-onset depression 3.6 (2.6–5.2)
Breslau et al. [15]	IHS migraine criteria	3.5 (2.6–4.6)	New-onset migraine 2.8 (2.2–3.5) New-onset depression 2.4 (1.8–3.0)
Breslau et al. [42]	IHS migraine criteria	Not assessed	New-onset migraine 3.4 (1.4–8.7) New-onset depression 5.8 (2.7–12.3)
Cross-sectional studies			
Merikangas [43]	Diagnosis by neurologist	2.2 (1.1–4.8)	Not assessed
Breslau et al. [15]	IHS migraine criteria	3.5 (2.6–4.6)	New-onset migraine 2.8 (2.2–3.5) New-onset depression 2.4 (1.8–3.0)
Swartz [44]	IHS migraine criteria	3.1 (2.0–4.4)	New-onset migraine 0.68 (0.02–2.0)
Zwart [45]	IHS migraine criteria	2.7 (2.3–3.2)	Not assessed
McWilliams [46]	Diagnosis by neurologist	2.8 (2.2–3.7)	Not assessed
Patel [47]	IHS migraine criteria	Strict migraine 2.7 (2.2–3.3) Probable migraine 1.9 (1.5–2.4)	Not assessed

Table 3 Studies of psychiatric comorbidity in patients with chronic daily headache

Study	Criteria for headache diagnosis	Criteria for psychiatric diagnosis	Mood disorders (%)	Anxiety disorders (%)
Mathew et al. [54]	By their own criteria	MMPI subscore	61	Not assessed
Verri et al. [10]	IHS	DSM IIIR (SCID-P)	Major depressive disorder: 25 dysthymia: 17	GAD: 69.3
Puca et al. [34]	IHS	CIDI-by physicians	45	56
Mitsikostas and Thomas [49]	IHS + mixed type headache	Psychiatric evaluation if HDRS > 16	10 (10 in CTTH, 19 in MOH, 9 in mixed type)	Not assessed
Wang et al. [76]	Adopted criteria of Silberstein et al. [2] + IHS for CTTH	DSM IV (GDS-S)	29	Not assessed
Juang et al. [48]	Adopted criteria of Silberstein et al. [2]	DSM IV (MINI)	Major depressive disorder: 55 dysthymia: 11	GAD: 5 Panic: 27
Holroyd et al. [41]	IHS	BDI	Major depressive disorder: 21	17
Atasoy et al. [75]	ICHD II	DSM IV (SCID-II, SCID-CV)	37.3 (25 in Group C; 51.6 in Group E; 36.2 in Group M)	15.3 (10.7 in Group C; 16.1 in Group E; 17.2 in Group M)

MINI International Neuropsychiatric Interview; *SCID* the Structured Clinical Interview for DSM-IV; *GDS-S* Geriatric Depression Scale-short Form; *IHS* International Headache Society; *CTTH* Chronic Tension-type Headache; *MMPI* Minnesota Multiphasic Personality Inventory; *HDRS* Hamilton Depression Rating Scale; *GAD* Generalized Anxiety Disorder; *BDI* Beck Depression Inventory; *CIDI* Composite International Diagnostic Interview; *ICHD-II* The international classification of headache disorders, 2nd edition; *MOH* Medication-Overuse Headache

study, high MMPI scores have been reported for the hypochondriasis and hysteria scales in CDH patients [55].

Galli and colleagues noted that only 9.6% of patients did not have familial occurrence of headache, with a half of the sample (49%) showing both parents and grandparents complaining of headache. Galli et al. [56] stressed the importance of familial recurrence (mainly for the mother) for migraine and psychiatric disorders, but less so for tension-type headaches and psychiatric disorders. The tension-type headache group had higher temperament traits of emotionality and shyness, and lower sociability, than the migraine group [57]. A higher emotionality score was also found among the parents and siblings of tension-type headache children [57]. Thus, high scores for emotionality and shyness can be considered to be symptoms of ‘behavioral inhibition’ [58, 59], characterized by the traits of irritability, shyness, fearfulness, and introversion. In school-age children, the presence of this temperament seems to increase the vulnerability to depressive and multiple anxiety disorders [60, 61]. Tension-type headache patients may have, as a group, more behavioral, emotional, and temperament difficulties than children referred for migraine [56]. This finding seems to be in contrast with an epidemiological study in Finland, which found that psychiatric symptoms tended to be more strongly associated with migraine than with tension-type headache [62]. However, personality disorders are considered to be a complication for headache management [63–65], and significant headaches are a complaint of about 60% of patients with

personality disorders presenting for acute treatment at a hospital emergency department [65].

Suicide risk

Suicide attempts seem to be more frequent in patients suffering from migraine than in the general population, especially in females and in patients with migraine with aura. This association has also been found in the general population [22, 35, 43, 66]. In contrast, the subtypes of CDH, headache frequencies, or medication overuse were not correlated with the risk of attempted suicide [66].

Migraine with aura also independently predicted a high suicidal risk (score > 10 on the MINI Suicidality Module) in adolescents with CDH [66], despite the fact that it is very difficult to predict suicide in youths. A history of migraine with aura, but not migraine without aura, has been found to be associated with an increased frequency of suicidal ideation and suicide attempts in patients with major depression in several studies [22, 39, 66], both current and previous affective episodes [20, 28]. Oedegaard and colleagues found that only 17% of patients having migraine aura without headache had made a suicide attempt, and these patients has the lowest frequency of affective temperaments as well as a higher age of onset of migraine auras [28]. However, the frequencies of suicidal ideation were approximately equal in the two groups of migraine patients [28].

Breslau [39] found an association of migraine with aura with suicide attempts that was not necessarily due to coexisting major depression. Migraine with aura alone was associated with significantly higher rates of suicidal tendencies. The risk of suicidal inclinations associated with migraine with aura and major depressive disorder was much higher than that for those with only major depressive disorder [67]. CDH is a source of great distress, with impairment of pleasure and working activities. This decreased quality of life can result in hopelessness and despair, and suicidal wishes can easily emerge [67]. However, the pain of headaches is itself a potential independent risk factor for suicide, particularly among those with chronic headache or multiple sources of co-occurring pain [68]. Individuals suffering from chronic pain may be particularly appropriate for suicide screening and intervention efforts. Innamorati and colleagues [69] have proposed a new scale, the Italian Perceived Disability Scale, as screening tool to identify comorbidity with emotional distress and disorders. This scale has proved able to predict suicidal intent in CDH patients and to assess disability in a sample of patients with CDH [69].

Treatment

The use of psychoactive drugs can influence the frequency and modify the clinical features of migraine headaches. For acute treatment, sumatriptan, nonsteroidal anti-inflammatory agents, ergotamine, and dihydroergotamine have been used, and metoclopramide is often co-administered.

The SSRIs (paroxetine and fluvoxamine) remain the first-choice medications for mood disorders related to migraine, although long-term treatment with serotonergic compounds does not impact the migraine itself. The SSRIs can also be used for the long-term treatment of panic disorder without mood disorders, tension-type headache, obsessive-compulsive disorder and somatoform disorder [70]. Regarding GAD and tension-type headache, the administration of BZDs is possible, but long-term therapy with BZD is not feasible because of the development of tolerance. In GAD long term therapy, SSRI (paroxetine) could represent a valid alternative to BZD if starting with a minimal effective dosage. The aim is to customize treatment as much as possible. The association between migraine and panic disorder and bipolar disorder or unipolar disorder can be treated with VPA, which is effective in the control of panic [71].

For the preventive treatment of mood disorders related to migraine or tension-type headache, the recommended therapy is to use antidepressive tricyclics (TCAs) [72], such as amitriptyline, as monotherapy or a combination of amitriptyline and an SSRI such as paroxetine [16] or flunarizine [73] and mood stabilizers such as lithium [21] or valproate [73, 74]. These have been tested and found to be

Table 4 Preventive treatments in headache associated with depressive syndrome

Preventive treatment
Migraine + depression + anxious symptoms: antidepressants with sedative properties
TCAs: amitriptyline
SSRIs: paroxetine, fluvoxamine
SNRIs: mirtazepine
Migraine + depression with psychomotor slowing: antidepressants without sedative properties
SSRIs: fluoxetine
Inadvisable drugs
Flunarizine
Beta-blockers

Table 5 Preventive treatments in headache associated with panic disorder

<i>Short-term treatment</i>
BZDs: alprazolam, clonazepam
<i>Long-term treatment</i>
Migraine + panic disorder: VPA
Migraine + panic disorder + bipolar disorder: VPA
Migraine + panic disorder + secondary depression: antidepressants with sedative properties
TCAs: amitriptyline
SSRIs: paroxetine, fluvoxamine
SNRIs: mirtazepine
Migraine + depression + secondary panic disorder
Prevalence of anxiety symptoms: antidepressants with sedative properties
TCAs: amitriptyline
SSRIs: paroxetine, fluvoxamine
SARI: trazodone
Prevalence of psychomotor slowing: antidepressants without sedative properties
SSRIs: fluoxetine

VPA Valproate

useful for migraine prophylaxis. No studies have reported a positive effect of the SSRIs on the aura symptoms. The better strategies are summarized in Tables 4 and 5.

Although, short-term pharmaceutical care intervention improves patients' mental health, it does not significantly change the number and severity of headaches. However, the increase in self-efficacy and mental health associated with pharmaceutical care may be instrumental in improving the long-term pharmacotherapy of patients with migraine and headache.

Conclusion

Several studies have analyzed the prevalence and impact of psychiatric disorders among headache patients, and these

studies have resulted in a better understanding of the relationship between migraine and mood disorders, anxiety, and suicide attempts. Unfortunately, as noted by Silberstein et al. [2], the relationship between headache and psychopathology has often been discussed only clinically rather than systematically studied. Furthermore, the investigation of the interplay of factors that precipitate suicide risk should include the assessment of chronic headache and its effects on well-being. These issues, as well as those discussed in this paper, represent an important area for future research.

Acknowledgments This Tutorial results partly from the teaching of the MP in the Master in Headache Medicine program, Sapienza University of Rome, during the academic year 2008–2009 (<http://w3.uniroma1.it/headache>). This academic educational activity is performed under the supervision of the WHO Global Campaign to Reduce the Burden of Headache Worldwide “Lifting The Burden”—Working Group for Specialist Education (SEWG).

Conflict of interest None.

References

- Dodick DW (2006) Chronic daily headache. *N Engl J Med* 354:158–165. doi:10.1056/NEJMc042897
- Silberstein SD, Lipton RB, Sliwinski M (1996) Classification of daily and near-daily headaches: field trial of revised HIS criteria. *Neurology* 47:871–875
- Scher AI, Stewart WF, Liberman J, Lipton RB (1998) Prevalence of frequent headache in a population sample. *Headache* 38:497–506. doi:10.1046/j.1526-4610.1998.3807497.x
- Lu SR, Fuh JL, Chen WT, Juang KD, Wang SJ (2001) Chronic daily headache in Taipei, Taiwan: prevalence, follow-up and outcome predictors. *Cephalalgia* 21:980–986. doi:10.1046/j.1468-2982.2001.00294.x
- Castillo J, Munoz P, Guitera V, Pascual J (1999) Epidemiology of chronic daily headache in the general population. *Headache* 39:190–196. doi:10.1046/j.1526-4610.1999.3903190.x
- Headache Classification Committee of the International Headache Society (2004) The international classification of headache disorders, 2nd edition. *Cephalalgia* 24(Suppl 1):9–160. doi:10.1111/j.1468-2982.2003.00824.x
- Silberstein SD, Olesen J, Bousser M, International Headache Society et al (2005) The international classification of headache disorders, 2nd edition (ICHD-II)—revision of criteria for 8.2 Medication-overuse headache. *Cephalalgia* 25:460–465. doi:10.1111/j.1468-2982.2005.00878.x
- Bigal ME, Tepper SJ, Sheftell FD et al (2004) Chronic daily headache: correlation between the 2004 and the 1988 international headache society diagnostic criteria. *Headache* 44:684–691. doi:10.1111/j.1526-4610.2004.04128.x
- Breslau N, Davis GC (1993) Migraine, physical health and psychiatric disorder: a prospective epidemiologic study in young adults. *J Psychiatr Res* 27:211–221. doi:10.1016/0022-3956(93)90009-Q
- Verri AP, Proietti Cecchini A, Galli C et al (1998) Psychiatric comorbidity in chronic daily headache. *Cephalalgia* 18(Suppl 21):45–49
- Mathew N (1991) Chronic daily headache: clinical features and natural history. In: Nappi G (ed) *Headache and depression: serotonin pathways as a common clue*. Raven Press, New York, pp 49–58
- Peres MF, Zukerman E, Young WB, Silberstein SD (2002) Fatigue in chronic migraine patients. *Cephalalgia* 22:720–724. doi:10.1046/j.1468-2982.2002.00426.x
- Peres MF, Young WB, Kaup AO et al (2001) Fibromyalgia is common in patients with transformed migraine. *Neurology* 57:1326–1328
- Puca F, the Italian Collaborative Group (2000) Psychological and social stressors and psychiatric comorbidity in patients with migraine without aura from headache centers in Italy a comparison with tension-type. *J Headache Pain* 1:17–32. doi:10.1007/s101940050005
- Breslau N, Schultz LR, Stewart WF et al (2000) Headache and major depression: is it the association specific to migraine? *Neurology* 54:308–313
- Sheftell FD, Atlas SJ (2002) Migraine and psychiatric comorbidity: from theory and hypotheses to clinical application. *Headache* 42:934–944. doi:10.1046/j.1526-4610.2002.02217.x
- Merikangas KR, Merikangas JR, Angst J (1993) Headache syndromes and psychiatric disorders: association and family transmission. *J Psychiatr Res* 2:197–210. doi:10.1016/0022-3956(93)90008-P
- Franchini L, Serretti A, Gasperini M, Smeraldi E (1998) Familial concordance of fluvoxamine response as a tool for differentiating mood disorder pedigrees. *J Psychiatr Res* 32:255–259. doi:10.1016/S0022-3956(98)00004-1
- Mahmood T, Silverstone T (2000) Twin concordance for bipolar disorder and migraines. *Am J Psychiatry* 157(12):2057. doi:10.1176/appi.ajp.157.12.2057
- Oedegaard KJ, Angst J, Neckelmann D, Fasmer OB (2005) Migraine aura without headache compared to migraine with aura in patients with affective disorders. *J Headache Pain* 6(5):378–386. doi:10.1007/s10194-005-0236-4
- Franchini L, Bongiorno F, Dotoli D et al (2004) Migraine headache and mood disorders: a descriptive study in an outpatient psychiatric population. *J Affect Disord* 81(2):157–160. doi:10.1016/S0165-0327(03)00164-2
- Breslau N, Davis GC, Andresky P (1991) Migraine, psychiatric disorders, and suicide attempts: an epidemiologic study of young adults. *J Psychiatr Res* 37:11–23. doi:10.1016/0165-1781(91)90102-U
- Post RM, Silberstein SD (1994) Shared mechanisms in affective illness, epilepsy, and migraine. *Neurology* 44(Suppl 7):S37–S47
- Pandey GN, Dwivedi Y, Rizavi HS et al (2002) Higher expression of serotonin 5-HT_{2A} receptors in the postmortem brains of teenage suicide victims. *Am J Psychiatry* 159:419–429. doi:10.1176/appi.ajp.159.3.419
- Lidberg L, Belfrage H, Bertilsson L et al (2000) Suicide attempts and impulse control disorder are related to low cerebrospinal fluid 5-HIAA in mentally disordered offenders. *Acta Psychiatr Scand* 101:395–402. doi:10.1034/j.1600-0447.2000.101005395.x
- Hargreaves RJ, Shephard SL (1999) Pathophysiology of migraine—new insights. *Can J Neurol Sci* 26(Suppl 3):S12–S19
- Shiah IS, Ko HC, Lee JF, Lu RB (1999) Platelet 5-HT and plasma MHPG levels in patients with bipolar I and bipolar II depressions and normal controls. *J Affect Disord* 52:101–110. doi:10.1016/S0165-0327(98)00060-3
- Fasmer OB, Oedegaard KJ (2001) Clinical characteristics of patients with major affective disorders and comorbid migraine. *World J Biol Psychiatry* 2:149–155
- Mongini F, Rota E, Deregibus A et al (2006) Accompanying symptoms and psychiatric comorbidity in migraine and tension-type headache patients. *J Psychosom Res* 61(4):447–451. doi:10.1016/j.jpsychores.2006.03.005

30. Abu-Arefeh I, Russell G (1995) Prevalence of headache and migraine in schoolchildren. *Br Med J* 309:765–769
31. Sillanpää M (1983) Prevalence of headache in prepuberty. *Headache* 23:10–14
32. Fasmer OB (2001) The prevalence of migraine in patients with bipolar and unipolar affective disorders. *Cephalalgia* 21(9):894–899. doi:10.1046/j.1468-2982.2001.00279.x
33. Endicott NA (1989) Psychophysiological correlates of “bipolarity”. *J Affect Disord* 17:47–56. doi:10.1016/0165-0327(89)90023-2
34. Puca F, the Italian Collaborative Group (1999) Psychiatric comorbidity and psychosocial stress in patients with tension-type headache from headache centers in Italy. *Cephalalgia* 19:159–164. doi:10.1046/j.1468-2982.1999.1903137.x
35. Merikangas KR, Stevens DE, Angst J (1993) Headache and personality: results of a community sample of young adults. *J Psychiatr Res* 27:187–196. doi:10.1016/0022-3956(93)90007-0
36. Merikangas KR, Stevens DE, Angst J (1994) Psychopathology and headache syndromes in the community. *Headache* 34:S17–S22. doi:10.1111/j.1526-4610.1994.hed3408s17.x
37. Breslau N, Merikangas K, Bowden CL (1994) Comorbidity of migraine and major affective disorders. *Neurology* 44(10 suppl 7): S17–S22
38. Breslau N, Andresky P (1995) Migraine, personality and psychiatric comorbidity. *Headache* 35:382–386. doi:10.1111/j.1526-4610.1995.hed3507382.x
39. Breslau N (1992) Migraine, suicidal ideation, and suicide attempts. *Neurology* 42:392–395
40. Merikangas KR, Stevens DE (1997) Comorbidity of migraine and psychiatric disorders. *Neurol Clin* 15:115–123. doi:10.1016/S0733-8619(05)70298-X
41. Holroyd KA, Stensland M, Lipchik GL et al (2000) Psychosocial correlates and impact of chronic tension-type headaches. *Headache* 40:3–16. doi:10.1046/j.1526-4610.2000.00001.x
42. Breslau N, Lipton RB, Stewart WF et al (2003) Comorbidity of migraine and depression: Investigating potential etiology and prognosis. *Neurology* 60:1308–1312
43. Merikangas KR, Angst J, Isler H (1990) Migraine and psychopathology: results of the Zurich cohort study of young adults. *Arch Gen Psychiatry* 47:849–853
44. Swartz KL, Pratt LA, Armenian HK et al (2000) Mental disorders and the incidence of migraine headaches in a community sample. *Arch Gen Psychiatry* 57:945–950. doi:10.1001/archpsyc.57.10.945
45. Zwart JA, Dyb G, Hagen K et al (2003) Depression and anxiety disorders associated with headache frequency. The Nord-Trøndelag Study. *Eur Neurol* 10:147–152. doi:10.1046/j.1468-1331.2003.00551.x
46. McWilliams LA, Goodwin RD, Cox BJ (2004) Depression and anxiety associated with three pain conditions: results from a nationally representative sample. *Pain* 111:77–83. doi:10.1016/j.pain.2004.06.002
47. Patel NV, Bigal ME, Kolodner KB et al (2004) Prevalence and impact of migraine and probable migraine in a health plan. *Neurology* 63:1432–1438
48. Juang KD, Wang SJ, Fuh JL et al (2000) Comorbidity of depressive and anxiety disorders in chronic daily headache and its subtypes. *Headache* 40:818–823. doi:10.1046/j.1526-4610.2000.00148.x
49. Mitsikostas DD, Thomas AM (1999) Comorbidity of headache and depressive disorders. *Cephalalgia* 19:211–217. doi:10.1046/j.1468-2982.1999.019004211.x
50. De Benedittis G, Lorenzetti A (1992) Minor stressful life events (daily hassles) in chronic primary headache relationship with MMPI personality patterns. *Headache* 32:330–332. doi:10.1111/j.1526-4610.1992.hed3207330.x
51. Fernandez E, Sheffield J (1996) Relative contributions of life events versus daily hassles to the frequency and intensity of headaches. *Headache* 36:595–602. doi:10.1046/j.1526-4610.1996.3610595.x
52. Marazziti D, Toni C, Pedri S, Bonuccelli U et al (1995) Headache, panic disorder and depression: comorbidity or a spectrum? *Neuropsychobiology* 31:125–129. doi:10.1159/000119182
53. Vasconcelos LP, Costa Silva M, Coelho Costa EA, da Silva AA Jr, Santiago Gomez R, Teixeira AL (2008) Obsessive compulsive disorder and migraine: case report, diagnosis and therapeutic approach. *J Headache Pain* 9:397–400
54. Mathew NT, Reuveni U, Perez F (1987) Transformed or evolutive migraine. *Headache* 27:102–106. doi:10.1111/j.1526-4610.1987.hed2702102.x
55. Mongini F, Defilippi N, Negro C (1997) Chronic daily headache. A clinical and psychological profile before and after treatment. *Headache* 37:83–87. doi:10.1046/j.1526-4610.1997.3702083.x
56. Galli F, Canzano L, Scalisi G, Guidetti V (2009) Psychiatric disorders and headache familial recurrence: a study on 200 children and their parents. *J Headache Pain* 10:187–197. doi:10.1007/s10194-009-0105-7
57. Mazzone L, Vitiello B, Incorpora G, Mazzone D (2005) Behavioural and temperamental characteristics of children and adolescents suffering from primary headache. *Cephalgia* 26:194–201. doi:10.1111/j.1468-2982.2005.01015.x
58. Garcia-Coll C, Kagan J, Reznick JS (1984) Behavioural inhibition in young children. *Child Dev* 55:1005–1019. doi:10.2307/1130152
59. Kagan J, Reznick JS, Clarke C et al (1984) Behavioural inhibition to the unfamiliar. *Child Dev* 55:2212–2225. doi:10.2307/1129793
60. Goodyer IM, Ashby L, Altham PME et al (1993) Temperament and major depression in 11–16 year olds. *J Child Psychol Psychiatry* 34:1409–1423. doi:10.1111/j.1469-7610.1993.tb02099.x
61. Kelvin RG, Goodyer IM, Altham PME (1996) Temperament and psychopathology amongst siblings of probands with depressive and anxiety disorders. *J Child Psychol Psychiatry* 37:543–550. doi:10.1111/j.1469-7610.1996.tb01440.x
62. Anttila P, Sourander A, Metsähonkala L, Aromaa M, Helenius H, Silanpää M (2004) Psychiatric symptoms in children with primary headache. *J Am Acad Adolesc Psychiatry* 43:412–419. doi:10.1097/00004583-200404000-00007
63. Saper JR, Lake AE III (2002) Borderline personality disorder and the chronic headache patient: review and management recommendations. *Headache* 42:663–674. doi:10.1046/j.1526-4610.2002.02156.x
64. Hegarty A (1993) The prevalence of migraine in borderline personality disorder. *Headache* 33:271
65. Stewart WF, Linet MS, Celentano DD (1989) Migraine headaches and panic attacks. *Psychosom Med* 51:559–569
66. Wang SJ, Juang KD, Fuh JL, Lu SR (2007) Psychiatric comorbidity and suicide risk in adolescents with chronic daily headache. *Neurology* 68(18):1468–1473. doi:10.1212/01.wnl.0000260607.90634.d6
67. De Filippis S, Erbutto D, Gentili F et al (2008) Mental turmoil, suicide risk, illness perception, and temperament, and their impact on quality of life in chronic daily headache. *J Headache Pain* 9:349–357. doi:10.1007/s10194-008-0072-4
68. Ilgen MA, Zivin K, McCammon RJ, Valenstein M (2008) Pain and suicidal thoughts, plans and attempts in the United States. *Gen Hosp Psychiatry* 30(6):521–527. doi:10.1016/j.genhosppsych.2008.09.003
69. Innamorati M, Pompili M, De Filippis S et al (2009) The validation of the Italian Perceived Disability Scale (IPDS) in chronic daily headache sufferers. *J Headache Pain* 10:21–26. doi:10.1007/s10194-008-0076-0

70. Savarese M, Guazzelli M, Prudenzeno MP et al (2005) Tertiary treatment for psychiatric comorbidity in headache patients. *J Headache Pain* 6(4):231–233. doi:[10.1007/s10194-005-0193-y](https://doi.org/10.1007/s10194-005-0193-y)
71. Malcangio M, Bowery NG (1996) GABA and its receptors in the spinal cord. *Trends Pharmacol Sci* 17:457–462. doi:[10.1016/S0165-6147\(96\)01013-9](https://doi.org/10.1016/S0165-6147(96)01013-9)
72. Singh NN, Misra S (2002) Sertraline in chronic tension-type headache. *J Assoc Physicians India* 50:871–872
73. Mitsikostas DD, Polychronidis I (1997) Valproate versus flunarizine in migraine prophylaxis: a randomized, double-open, clinical trial. *Fund Neurol* 12:267–276
74. Nieper HA (1973) Clinical applications of lithium orotate: a two year study. *J Acad Prevent Med* 2:18–22
75. Atasoy HT, Atasoy N, Unal AE, Emre U, Sumer M (2005) Psychiatric comorbidity in medication overuse headache patients with pre-existing headache type of episodic tension-type headache. *Eur J Pain* 9(3):285–291
76. Wang SJ, Fuh JL, Lu SR, Liu CY, Hsu LC, Wang PN, Liu HC (2000) Chronic daily headache in Chinese elderly: prevalence, risk factors, and biannual follow-up. *Neurology* 54:314–319